

Geometry of Quantum Group Twists, Multidimensional Jackson Calculus and Regularization

A.P.Demichev[†]

Centro Brasileiro de Pesquisas Fisicas - CBPF/CNPq,
Rua Dr.Xavier Sigaund, 150,
22290-180, Rio de Janeiro,
RJ Brasil

Abstract

We show that R-matrices of all simple quantum groups have the properties which permit to present quantum group twists as transitions to other coordinate frames on quantum spaces. This implies physical equivalence of field theories invariant with respect to q-groups (considered as q-deformed space-time groups of transformations) connected with each other by the twists. Taking into account this freedom we study quantum spaces of the special type: with commuting coordinates but with q-deformed differential calculus and construct $GL_r(N)$ invariant multidimensional Jackson derivatives. We consider a particle and field theory on a two-dimensional q-space of this kind and come to the conclusion that only one (time-like) coordinate proved to be discretized.

[†]on leave of absence from Nuclear Physics Institute, Moscow State University, 119899, Moscow, Russia

1 Introduction

Lattice regularization has many advantages and plays important role in quantum field theory (see e.g. [1]). Unfortunately, it has some shortcomings also. Perhaps the most essential one is a space-time symmetry breaking. The general reason for the latter is connected with introduction of a lattice in a theory *by hands* and so with its too rigid nature. Among other reasons this fact initiated many attempts to construct discrete ("quantized") space-time manifolds on a more deep background (about old attempts see e.g. [2], ch. VII and refs. therein). Last years this problem has received revival and considerable interest due to appearance of quantum groups [3], [4], [5].

Quantum spaces which appear in the frame of quantum group theory [5] have many unusual properties, in particular, q -deformed differential calculi [6] and, in general, non-commuting coordinates. In one-dimensional space q -derivative can be represented by Jackson difference operator [7],[8]. This, in turn, provides description of a quantum particle on a one-dimensional lattice [9]. Thus a q -deformation of a differential calculus apparently leads to space discretization. The relation of the non-commutativity of coordinates to space discretization is not so straightforward. From the other hand, the non-commutativity causes many problems in construction of field theories on q -spaces. Indeed, it means that operators of coordinates can not be diagonalized and have not definite values simultaneously. This essentially new property of q -spaces is not disastrous for the approach which aims to construct new and more fundamental theory of space-time, gravity, etc. An interesting attempts in this direction was made in [10]. From this work it is clear also that it is special property of phase subspace for each pair of coordinate and conjugate momentum that leads to improved ultraviolet behaviour of the model (perturbation theory is based on elementary excitations with fast growing energy spectrum of q -oscillators, roughly $E_n \sim [n; q] \equiv (q^n - 1)/(q - 1)$, where $q > 1$). The non-commutativity looks like "payment" for existence of some q -deformed symmetry ($SU_q(N)$ in this case, N being space-time dimensionality). As is seen from the paper, the physical properties of the model are quite unusual. In particular, there are no states with definite values of position or momentum coordinates and even with definite values of any part of them. So the very notion of a particle becomes ill defined. This means that models of this kind can not be considered as a regularization of usual quantum field theory but prove to be of essentially new type.

These properties of q -deformed models seems to cause serious problems if one aims to construct in this way lattice-like regularization of *existent* field theories. It is obvious that the necessary condition for the such regularization (which does not bring essentially new properties to field theoretical models on a continuous space-time) is a commutativity of coordinates. This letter is devoted to the studying of such possibility. The key idea of our approach is a physical equivalence of different q -deformations of space-time manifolds which are related to each other by twists of corresponding quantum groups. As was shown in [11], q -deformed Minkowski space-time with non-

commuting coordinates which corresponds to pure twisted Poincaré group (i.e. to the q-group obtained from the classical one by a twist) can be constructed from a usual Minkowski space with help of appropriate coordinate transformation and q-generalization of 4-beins. In this letter we generalize this partial result to twists of non-trivially deformed all simple groups. More precisely, we will show that known R-matrices for all simple q-groups have the property which permits to describe the twist procedure as a transformation of q-space coordinates.

To construct lattice-like regularization one definitely needs multidimensional finite difference calculus. Using above mentioned freedom in choice of different q-spaces we consider $GL_q(N)$ -invariant q-spaces with commuting coordinates and q-deformed differential calculus and construct multidimensional analog of the Jackson calculus (invariant with respect to appropriate quantum group).

Using explicit formulas for a two-dimensional space we consider a quantum mechanical particle and simplest field theory and show that the latter is equivalent to the system on a cylinder with time coordinate taking values on equidistant lattice along the cylinder. Surprisingly, the second coordinate (space-like coordinate on a circle) proved to be continuous. This fact takes its origin in the properties of involution of the corresponding quantum general linear group which lead to construction of the quantum mechanical states in a mixed coordinate-momentum representation, so that "the second discreteness" corresponds to integer numbers labeling Fourier modes on the circles. Such discretization is enough for the regularization of two-dimensional theories. But this fact is dangerous for higher dimensional spaces.

2 Geometry of quantum group twists

As is shown in [12],[13], multiparametric quantum groups can be obtained from a one-parametric q-group via so called twists of a quasitriangular Hopf algebra A with help of an element $\mathcal{F} = \sum f^i \otimes f_i \in A \otimes A$, which satisfies certain relations, so that new coproduct $\Delta^{(F)}$ and new universal R-matrix $\mathcal{R}^{(F)}$ are connected with the initial objects Δ and $\mathcal{R}$ through the relations

$$\Delta^{(F)} = \mathcal{F} \Delta \mathcal{F}^{-1}, \quad \mathcal{R}^{(F)} = \mathcal{F}^{-1} \mathcal{R} \mathcal{F}^{-1} .$$

Consider at first the case of q-deformations of $GL(N)$ groups. In this case a twist of R-matrix in fundamental representation R is described with help of diagonal matrix $F = \text{diag}(f_{11}, f_{12}, \dots, f_{nn})$ with $f_{ij} f_{ji} = 1$ so that R-matrix $R^{(F)}$ of the twisted group $GL_{r, \tilde{q}_{ij}}(N)$ has a form

$$R^{(F)} = F^{-1} R F^{-1} .$$

Here R is (in general, also multiparametric) R-matrix of the initial group $GL_{r, q_{ij}}(N)$ and

$$\tilde{q}_{ij} = q_{ij} f_{ij}^2 . \tag{1}$$

Coordinates of the initial quantum space $C_q^N[x^i]$ satisfy the commutation relations (CR) [5],[14]

$$x^i x^j = q_{ij} x^j x^i \quad (2)$$

and coordinates of the twisted space $C_q^{(F)N}[\tilde{x}^i]$ have the CR

$$\tilde{x}^i \tilde{x}^j = \tilde{q}_{ij} \tilde{x}^j \tilde{x}^i . \quad (3)$$

Now we introduce the algebra $E_q^N[e^i, g_j]$ with the generators $\{e^i, g_i\}_{i=1}^N$ which commute with coordinates and put

$$\tilde{x}^i = e^i x^i \quad (\text{no summation}) . \quad (4)$$

The elements e^i play the role of components of a q-deformed (diagonal) N-bein. CR for them follows from (1)-(4)

$$e^i e^j = f_{ij}^2 e^j e^i , \quad (5)$$

and g_i are inverse elements

$$g_i e^i = 1 . \quad (6)$$

The coordinates $\tilde{x}^i$ are transformed by a q-matrix $\tilde{T}$

$$\tilde{x}'^i = \sum_{j=1}^N \tilde{T}_j^i \tilde{x}^j . \quad (7)$$

Then using (4),(6) one obtains from (7) transformations of the coordinates x^i

$$x'^i = \sum_{j=1}^N g_i \otimes \tilde{T}_j^i \otimes e^j \otimes x^j . \quad (8)$$

We used in (8) a cross-product sign to stress that the elements from the different sets commute with each other (the elements g_i in (8) must be considered as the inverse elements to generators e^i of another copy of an algebra E_q^N with respect to the elements e^i entering the same formula). The relation (8) means that the coordinates x^i are transformed by the matrix T with the entries

$$T_j^i = g_i \otimes \tilde{T}_j^i \otimes e^j \quad (\text{no summation}) . \quad (9)$$

Using (6) one can express the matrix $\tilde{T}$ through T

$$\tilde{T}_j^i = e^i \otimes T_j^i \otimes g_j \quad (\text{no summation}) . \quad (10)$$

One can check straightforwardly that x'^i defined by (8) satisfy the correct CR

$$x'^i x'^j = q_{ij} x'^j x'^i .$$

The general reason for this is the following property of the R-matrices: if a q-matrix T satisfies TT-relation defined by the corresponding R-matrix, then the $\tilde{T}$ -matrix defined by (9) or (10) satisfies the relation with twisted R-matrix $R^{(F)}$.

To prove this statement let us write TT-relation (CR for entries of a matrix T) in explicit form

$$\sum_{p,s} R_{ps}^{mn} T_u^p T_v^s = \sum_{s,r} T_s^n T_r^m R_{uv}^{rs}$$

and substitute T_j^i by their expressions (9) in terms of $\tilde{T}_j^i$. This gives the relation for the latters

$$\sum_{p,s} R_{ps}^{mn} g_p g_s e^u e^v \tilde{T}_u^p \tilde{T}_v^s = \sum_{s,r} \tilde{T}_s^n \tilde{T}_r^m g_n g_m e^s e^r R_{uv}^{rs} . \quad (11)$$

Remind that in this relation the elements g_i must be considered as inverse elements for the generators e^i of another copy of an algebra E_q^N and so they commute with the elements e^i entering the same relation.

Multiparametric R-matrix for $GL_{r,q_{ij}}(N)$ group has a form

$$R_{ps}^{mn} = B_{ps}^{mn} + N_{ps}^{mn} , \quad (12)$$

where B is the diagonal matrix

$$B_{ps}^{mn} = \delta_p^m \delta_s^n (\delta^{mn} + \Theta^{nm} q_{mn}^{-1} + \Theta^{mn} q_{nm} r^{-1}) , \quad (13)$$

with $\Theta^{mn} = 1$ if $m > n$, $\Theta^{mn} = 0$ if $m \leq n$, and the matrix N is off-diagonal part of the R-matrix

$$N_{ps}^{mn} = \delta_s^m \delta_p^n \Theta^{mn} (1 - r^{-1}) . \quad (14)$$

Using this expressions one easily obtains

$$B_{ps}^{mn} g_p g_s = g_m g_n B_{ps}^{mn} = f_{mn}^{-2} g_n g_m B_{ps}^{mn} = g_n g_m B_{ps}^{(F)mn} , \quad (15)$$

$$N_{ps}^{mn} g_p g_s = g_n g_m N_{ps}^{mn} , \quad (16)$$

so that

$$R_{ps}^{mn} g_p g_s e^u e^v = g_n g_m e^u e^v R_{ps}^{(F)mn} ,$$

where $R^{(F)}, B^{(F)}$ are the twisted matrices of the same form (12)-(14) but for the twisted parameters $\tilde{q}_{ij} = q_{ij} f_{ij}^2$. Analogous consideration for the RHS of (11) shows that this relation can be rewritten in the form

$$\sum_{p,s} R_{ps}^{(F)mn} \tilde{T}_u^p \tilde{T}_v^s = \sum_{s,r} \tilde{T}_s^n \tilde{T}_r^m R_{uv}^{(F)rs} . \quad \square \quad (17)$$

Thus twisted q-matrices can be constructed with the help of q-deformed N-beins (5),(6) and the formula (9) which is direct generalization (q-deformation) of a relation between matrices of transformations in different coordinate frames.

In the case of q-deformation of simple groups of the series B_N , C_N , D_N there is one more structure, namely an invariant length [5]

$$L_q = \sum_{i,j} x^i C_{ij} x^j = \sum_i l_i x^{i'} x^i ,$$

where $i' = N + 1 - i$. Values of the coefficients l_i can be found in [5] and are not essential for our consideration. To preserve L_q , components of a q-bein must satisfy the additional constraints

$$e^i e^{i'} = e^{i'} e^i = 1 , \quad (18)$$

$i = 1, \dots, N/2$ for C_N, D_N series; $i = 1, \dots, (N + 1)/2$ for B_N series. In particular, for the series B_N

$$e^{(N+1)/2} = 1 . \quad (19)$$

These constraints reduce number of twist parameters, which from the geometrical point of view define CR for the components of the q-beins, so that the number is equal to $k(k - 1)/2$, where k is rank of a group. R-matrices for the B_N , C_N , D_N series have the form

$$R_{kl}^{ij} = \left[\delta_k^i \delta_l^j \left(r \delta^{ij} (1 - \delta^{ii'}) + (\Theta^{ji} r q_{ij}^{-1} + \Theta^{ij} q_{ji} r^{-1}) (1 - \delta^{ii'}) \right) + (r - r^{-1}) \delta_l^i \delta_k^j \Theta^{ij} \right] +$$

$$+ \left[\frac{1}{r} \delta_k^i \delta_l^j \delta^{jj'} (1 - \delta^{ii'}) - \Theta^{ij} (r - r^{-1}) r^{(\rho_i - \rho_j)} \epsilon_i \epsilon_j \delta^{ij'} \delta_{kl'} + \delta_{(N+1)/2}^i \delta_{(N+1)/2}^j \delta^{(N+1)/2}_k \delta^{(N+1)/2}_l \right]$$

(the last term exists for the B_N series only). Here ρ_i and ϵ_i are integer or half integer numbers [5],[16]. Using this explicit form one can easily show that the R-matrices have the property analogous to that of the A_N groups. Indeed, the terms in the first square brackets have the structure similar to those of the R-matrix for the $GL_{r,q_{ij}}(N)$ groups. So they are transformed properly when the elements e_i, g_j move through them (cf. (15),(16)). The terms in the second square brackets are not changed because of Kronecker symbols $\delta^{jj'}$, $\delta^{ij'}$ or $\delta^{(N+1)/2}_k$, $\delta_{(N+1)/2}^i$ and the relations (18),(19).

Thus again the matrices $\tilde{T}$ defined by (9),(10) satisfy the CR (17) for twisted quantum groups.

3 q-Spaces with commuting coordinates and multidimensional Jackson differential calculus

It is well known that in one-dimensional case q-deformed differential calculus can be realized with help of finite difference operation called Jackson derivative [7],[8] which has the form

$$D_r f(x) = \frac{f(x) - f(rx)}{(1 - r)x} , \quad (20)$$

or

$$\tilde{D}_r f(x) = \frac{f(r^{-1}x) - f(x)}{(r^{-1} - 1)x} \quad (21)$$

with the CR

$$D_r x - r x D_r = 1, \quad \tilde{D}_r x - r^{-1} x \tilde{D}_r = 1$$

(throughout the paper we will assume that $r \leq 1$). In particular, these derivatives are suitable for description of a quantum mechanical particle on a one-dimensional lattice [9]. To consider quantum mechanics on higher dimensional lattices one needs multidimensional generalization of the Jackson calculus. Such calculus invariant with respect to the q-group $GL_r(N) := GL_{r,1}(N)$ can be constructed in the space $C_r^N[x^i]$ with commuting coordinates x^i .

As is shown in the previous section, commuting coordinates differ from non-commuting ones by the non-commuting factors e^i . The situation reminds a transition from usual three-dimensional Euclidian coordinates to well known quaternions with basis $\{\sigma_i\}_{i=1}^3$: sometimes it is convenient to put in the coorespondence to the coordinates $\{x^i\}_{i=1}^3$ non-commutative quaternions $\hat{x}^i := x^i \sigma_i$ (no summation). Though this transformation brings new algebraic structure and permits to express three-dimensional rotations in pure algebraic way, the underling geometrical and physical structure remains the same.

This analogy leads to the conclusion that one can freely choose a most convenient quantum space among a set of twisted q-spaces. In particular, in the case of spaces $C_{r,q_{ij}}^N[x^i]$ the most simple choice is the spaces $C_r^N[x^i]$ with commuting coordinates. The CR for coordinates and derivatives on this space are the following [14]

$$\begin{aligned} x^i x^j &= x^j x^i \quad \forall i, j, \quad \partial_i x^i = 1 + r x^i \partial_i + (r - 1) \sum_{l=i+1}^N x^l \partial_l, \\ \partial_i \partial_k &= \frac{1}{r} \partial_k \partial_i, \quad \partial_i x^k = r x^k \partial_i, \quad \partial_k x^i = x^i \partial_k, \\ &i < k, \quad i, k = 1, \dots, N. \end{aligned} \quad (22)$$

To develop Jackson calculus define the operators of finite dilatations

$$A_i = 1 + (r - 1) \sum_{j=i}^N x^j \partial_j \quad (23)$$

with the commutation relations

$$\begin{aligned} A_i A_k &= A_k A_i, \quad \forall k, i, \\ A_k x^i &= x^i A_k, \quad k > i, \\ A_i x^k &= r x^k A_i, \quad k \geq i. \end{aligned} \quad (24)$$

Note that the operators A_i are analogous to operators Y_j^i of vector fields on a simple quantum group, introduced in [15]. The relations (23) permit to express the q-derivatives in terms of A_i

$$\begin{aligned}\partial_i &= (1-r)^{-1}(x^i)^{-1}(A_{i+1} - A_i) , \quad i = 1, \dots, N \\ A_{N+1} &:= 1 .\end{aligned}\tag{25}$$

The relations (24) and (25) lead to the following realization of the q-derivatives in a space of functions of N commuting variables

$$\partial_i f(x^1, \dots, x^N) = \frac{f(x^1, \dots, x^i, rx^{i+1}, \dots, rx^N) - f(x^1, \dots, x^{i-1}, rx^i, \dots, rx^N)}{(1-r)x^i} .\tag{26}$$

One can check easily that the finite differences (26) satisfy the $GL_r(N)$ invariant CR (22) indeed. These differences look like natural multidimensional generalization of the Jackson derivative (20).

As is shown in [6] there are two types of CR for q-derivatives and coordinates, which are invariant with respect to q-deformed groups. The first possibility is presented in (22), the second one is the following

$$\begin{aligned}\tilde{\partial}_i x^i &= 1 + \frac{1}{r} x^i \tilde{\partial}_i + \left(\frac{1}{r} - 1\right) \sum_{l=1}^{i-1} x^l \tilde{\partial}_l , \\ \tilde{\partial}_i \tilde{\partial}_k &= \frac{1}{r} \tilde{\partial}_k \tilde{\partial}_i , \quad \tilde{\partial}_i x^k = x^k \tilde{\partial}_i , \quad \tilde{\partial}_k x^i = \frac{1}{r} x^i \tilde{\partial}_k , \\ &\quad i < k, \quad i, k = 1, \dots, N .\end{aligned}\tag{27}$$

For these CR it is natural to introduce the operators

$$\tilde{A}_i = 1 + \left(\frac{1}{r} - 1\right) \sum_{j=1}^i x^j \tilde{\partial}_j ,\tag{28}$$

which commute with each other and have the following CR with the coordinates

$$\begin{aligned}\tilde{A}_k x^i &= r^{-1} x^i \tilde{A}_k , \quad k > i , \\ \tilde{A}_i x^k &= x^k \tilde{A}_i , \quad k \geq i .\end{aligned}\tag{29}$$

These relations permit to construct the realization of the q-derivatives $\tilde{\partial}_i$ in terms of the finite differences

$$\tilde{\partial}_i f(x^1, \dots, x^N) = \frac{f(r^{-1}x^1, \dots, r^{-1}x^i, x^{i+1}, \dots, x^N) - f(r^{-1}x^1, \dots, r^{-1}x^{i-1}, rx^i, \dots, rx^N)}{(r^{-1} - 1)x^i} ,\tag{30}$$

which is the generalization of the Jackson derivative of the second type (21).

4 Quantum particle on two-dimensionall quantum space

In this section we apply the above considered formulas for construction of quantum mechanics on a two-dimensional quantum plane. For convenience we rewrite the CR (22), (24), (27) and (29) in this particular case denoting: $x^1 = z$, $x^2 = \bar{z}$, $\partial_1 = \partial$, $\partial_2 = \bar{\partial}$, $\tilde{\partial}_1 = \tilde{\partial}$, $\tilde{\partial}_2 = \tilde{\bar{\partial}}$, $A_1 = A$, $A_2 = \bar{A}$, $\tilde{A}_1 = \tilde{A}$, $\tilde{A}_2 = \tilde{\bar{A}}$

$$z\bar{z} = \bar{z}z, \quad \partial\bar{\partial} = \frac{1}{r}\bar{\partial}\partial, \quad \tilde{\partial}\tilde{\bar{\partial}} = \frac{1}{r}\tilde{\bar{\partial}}\tilde{\partial}, \quad (31)$$

$$\begin{aligned} \partial z &= 1 + rz\partial + (r-1)\bar{z}\bar{\partial}, & \bar{\partial}\bar{z} &= 1 + r\bar{z}\bar{\partial}, \\ \partial\bar{z} &= r\bar{z}\bar{\partial}, & \bar{\partial}z &= z\bar{\partial}, \end{aligned} \quad (32)$$

$$Az = rzA, \quad A\bar{z} = r\bar{z}A, \quad \bar{A}z = z\bar{A}, \quad \bar{A}\bar{z} = r\bar{z}\bar{A}, \quad (33)$$

$$\begin{aligned} \tilde{\partial}z &= 1 + \frac{1}{r}z\tilde{\partial}, & \tilde{\bar{\partial}}\bar{z} &= 1 + \left(\frac{1}{r} - 1\right)z\tilde{\partial} + \frac{1}{r}\bar{z}\tilde{\bar{\partial}}, \\ \tilde{\bar{\partial}}\bar{z} &= \bar{z}\tilde{\bar{\partial}}, & \tilde{\bar{\partial}}z &= \frac{1}{r}z\tilde{\bar{\partial}}, \end{aligned} \quad (34)$$

$$\tilde{A}z = \frac{1}{r}z\tilde{A}, \quad \tilde{A}\bar{z} = \bar{z}\tilde{A}, \quad \tilde{\bar{A}}z = \frac{1}{r}z\tilde{\bar{A}}, \quad \tilde{\bar{A}}\bar{z} = \frac{1}{r}\bar{z}\tilde{\bar{A}}, \quad (35)$$

All A-operators commute with each other and satisfy the relations

$$\tilde{\bar{A}}A = 1, \quad \tilde{\bar{A}}\bar{A} = \tilde{A}, \quad \tilde{A}A = \bar{A}. \quad (36)$$

The simplest way to find them is to derive from an action of the operators on an arbitrary function $f(z, \bar{z})$. The CR between different types of the q-derivatives have the form

$$\begin{aligned} \tilde{\partial}\partial &= r\partial\tilde{\partial}, & \tilde{\bar{\partial}}\bar{\partial} &= r\bar{\partial}\tilde{\bar{\partial}}, \\ \tilde{\bar{\partial}}\partial &= \bar{\partial}\tilde{\partial}, & \tilde{\bar{\partial}}\bar{\partial} &= r^2\bar{\partial}\tilde{\bar{\partial}}. \end{aligned} \quad (37)$$

Now we must define *-involution in the algebra of the operators which enter the relations (31)-(37).

We want to consider the parameter r as a lattice spacing, hence, it must be a real number. The appropriate involution for $GL_r(2)$ in this case is the following [16]

$$T^* = CTC, \quad C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

This means that

$$T^* = \begin{pmatrix} a^* & b^* \\ c^* & d^* \end{pmatrix} = \begin{pmatrix} d & c \\ b & a \end{pmatrix}$$

and $z^* = \bar{z}$. It is not difficult to see that the CR for the A-operators and the coordinates are consistent with the involution

$$A^* = \tilde{\bar{A}} , \quad \bar{A}^* = \tilde{A} . \quad (38)$$

This gives the involution rules for the derivatives

$$\partial^* = -\tilde{\bar{\partial}} + \left(\frac{1}{r} - 1\right) \frac{z}{\tilde{z}} \tilde{\partial} + \frac{1}{\tilde{z}} , \quad \bar{\partial}^* = -\tilde{\partial} + \frac{1}{z} .$$

These rules look rather combersome, but they are direct generalization of the involution in one-dimensional case [9].

The next step is to construct the representation of the operators in a Hilbert space so that the involution would coincide with hermitian conjugation (therefore in the following we will denote the involution by a sign of hermitian conjugation). To construct convenient basis of a Hilbert space one needs hermitian operators. Hermitian combinations of the coordinates of the form

$$x = (z + \bar{z})/2 , \quad y = (z - \bar{z})/2i$$

are not convenient as they have identical CR with a part of A-operators and rather combersome CR with another part. A-operators play the role of conjugate momenta on a lattice (cf. [9]). A natural choice of position operators follows from the observation that the A-operators generate finite dilatations rather than translations. This implies the introduction of the hermitian operator

$$\rho = \sqrt{\tilde{z}z} , \quad \rho^\dagger = \rho , \quad (39)$$

and the unitary one

$$\Phi = \sqrt{\tilde{z}z^{-1}} , \quad \Phi^\dagger \Phi = 1 . \quad (40)$$

As follows from (38) and (36) we have also unitary operator A and hermitian operator $B := \tilde{A}\bar{A}$. These operators have the following CR

$$\begin{aligned} A\rho &= r\rho A , & B\rho &= \rho B , \\ A\Phi &= \Phi A , & B\Phi &= r\Phi B . \end{aligned}$$

All other operators can be expressed in terms of these four ones. Thus we have two commuting hermitian operators ρ , B and can define a basis $|N, m\rangle$ in a Hilbert space $\mathcal{H}$ in which these operators are diagonal (cf. [9], [17])

$$\begin{aligned} \rho |N, m\rangle &= \rho_0 r^{-N} |N, m\rangle \\ B |N, m\rangle &= b_0 r^{-m} |N, m\rangle . \end{aligned} \quad (41)$$

The constants ρ_0 and b_0 mark different equivalent representations and for shortness we will put $\rho_0 = b_0 = 1$. The unitary operators transform a vector with given eigenvalue into the one with a neighboring eigenvalue

$$\begin{aligned} A |N, m\rangle &= |N+1, m\rangle \\ \Phi |N, m\rangle &= |N, m-1\rangle . \end{aligned} \quad (42)$$

So starting from $GL_r(2)$ invariant differential calculus on a quantum plane we have come naturally to the polar coordinates, the operator ρ being the operator of radial coordinates and the operator Φ being of the form $\Phi = e^{-i\phi}$, where ϕ is (hermitian) operator of an angle coordinate. The structure of the algebra involution leads to the mixed representation with one coordinate (ρ) and one momentum diagonal operators. In classical case a multiplication of a function by Φ shifts its Fourier components numbers by minus unity. This corresponds to the action (42) of the operator Φ on vectors of $\mathcal{H}$. This implies that B is connected with an angular momentum operator. In fact it is analog of the operator of the form

$$\exp\{i\phi_0 \frac{\partial}{\partial \phi}\} = \exp\{\phi_0 M\} \quad (43)$$

in the case of quantum mechanics on usual continuous plane, where ϕ_0 is some fixed angle and $i\partial/\partial\phi$ is (two-dimensional) angular momentum operator. Eigenfunctions of this operator are periodic functions $\exp\{in\phi\}$ with eigenvalues $\exp\{n\phi_0\} = (e^{\phi_0})^n$. The latter expression coincides with eigenvalue of operator B if one equates $r = \exp\{\phi_0\}$.

Thus the operator ρ defines values of radial coordinate and operator the A shifts them (play the role of conjugate momentum). Analogously, the operator B defines angular momentum values and the operator Φ of the conjugate coordinate shifts its eigenvalues.

Now we can consider a q-subgroup Λ of the $GL_r(2, \mathbb{R})$ of matrices of the form

$$T = \begin{pmatrix} a & 0 \\ 0 & a^* \end{pmatrix},$$

with $aa^* = a^*a$. Because of the latter relation it looks like ordinary group isomorphic to a multiplicative group of complex numbers. But one must remember about CR with the generators Y_j^i of the corresponding quantum universal enveloping algebra. For left-invariant and right-covariant generators they have the following general form [15]

$$Y_j^i T_s^k = T_l^k Y_n^m (\tilde{R}_{21})_{ml}^{il} (\tilde{R}_{12})_{js}^{nt}, \quad (44)$$

where $\tilde{R}_{12}$ and $\tilde{R}_{21}$ are properly normalized R-matrices of the $GL_r(2, \mathbb{R})$. The CR (44) gives for the subgroup Λ (i.e. if $b = c = 0$)

$$\begin{aligned} Y_1^1 a &= ra Y_1^1, & Y_1^1 a^* &= a^* Y_1^1, \\ Y_2^2 a &= a Y_2^2, & Y_2^2 a^* &= ra^* Y_2^2. \end{aligned} \quad (45)$$

It is easy to see that CR for $\mathcal{D} := Y_1^1 Y_2^2$, $\bar{\mathcal{D}} := Y_2^2$, a and a^* are the same as for the $A, \tilde{A}, z, \bar{z}$. So the algebra (33) on the quantum plane is isomorphic to that of the q-subgroup Λ and we can identify the q-plane with this subgroup. The operators $\tilde{A}, \bar{\tilde{A}}$ correspond to right-invariant and left-covariant generators (cf. [15]). From the

other hand, q-subgroup Λ can play the role of symmetry group, the whole $GL_r(2, \mathbb{R})$ group being the group of linear canonical transformations (of CR (32),(34)).

The coaction of Λ on the coordinates is

$$\begin{aligned} z &\rightarrow z' = a \otimes z , & \bar{z} &\rightarrow \bar{z}' = a^* \otimes \bar{z} , \\ \rho &\rightarrow \rho' = a_\rho \otimes \rho , & \Phi &\rightarrow \Phi' = a_\Phi \otimes \Phi , \\ a_\rho &= \sqrt{a^* a} , & a_\Phi &= \sqrt{a/a^*} . \end{aligned} \quad (46)$$

The coordinates ρ' , Φ' have a representation in a Hilbert space $\mathcal{H} \otimes \mathcal{H}$ (cross product of the same Hilbert space as in our case the comodule coincides with the symmetry q-group) which has a subspace (diagonal) isomorphic to $\mathcal{H}$. The latter corresponds to the representation of CR based on the coordinates ρ' , Φ' as a primary ones. All A-operators and so the operator B are invariant with respect to the coaction (46).

To consider a two dimensional quantum free particle one can use, for example, the Hamiltonian

$$h = \frac{\Omega}{(1-r)^2} \left[(1-A)(1-A^\dagger) + (1-B)^2 \right] , \quad (47)$$

where Ω is constant with dimension of energy. In order to find eigenfunctions and eigenvalues of the Hamiltonian introduce the operators

$$\rho^{iP} = \exp\{iP \ln \rho\} .$$

and the states (cf. [17])

$$\begin{aligned} | \mathbb{1}, m \rangle &= \sum_{m=-\infty}^{\infty} | N, m \rangle , \\ | P, m \rangle &= \rho^{iP} | \mathbb{1}, m \rangle \end{aligned}$$

with the properties

$$\begin{aligned} A | \mathbb{1}, m \rangle &= | \mathbb{1}, m \rangle , \\ A | P, m \rangle &= r^{iP} | P, m \rangle , \end{aligned}$$

where P is a real number: $0 \leq P \leq \pi/\chi$, $\chi := \ln r$. For eigenvalues of the Hamiltonian (47) we obtain

$$\begin{aligned} h | P, m \rangle &= \xi_{P,m} | P, m \rangle , \\ \xi_{P,m} &= \Omega \left(\frac{(1 - \cos \chi P)}{(1-r)^2} + [m; r]^2 \right) \xrightarrow{r \rightarrow 1} \\ &\xrightarrow{r \rightarrow 1} \Omega(P^2 + m^2) , \quad [m; r] = \frac{1 - r^m}{1 - r} . \end{aligned}$$

Thus the operator h has the correct continuous limit but its eigenvalues are not invariant with respect to the reflection $m \rightarrow -m$. This means that left and right modes have different properties and positive modes have decreasing spectrum which

can lead to additional divergencies in corresponding field theories. These properties are caused by exponent-like form of the operator B analogous to (43) as we discussed above. So it is natural to consider the Hamiltonian

$$H = \frac{\Omega}{(1-r)^2} [(1-A)(1-A^\dagger) + \ln^2 B]$$

with the eigenvalues $\Omega\lambda_{P,m}$, where

$$\lambda_{P,m} = \left(\frac{(1 - \cos \chi^P)}{(1-r)^2} + \frac{\chi^2}{(1-r)^2} m^2 \right) \xrightarrow{r \rightarrow 1} (P^2 + m^2) . \quad (48)$$

To construct two dimensional quantum field theory we need a kind of integral over the variable ρ . It can be defined analogously to that in [9], [17].

For m-th component $f_m(\rho)$ of "Fourier expansion" we define

$$\begin{aligned} \int_0^K d_r \rho f_m(\rho) &:= \langle K, m | \partial_\rho^{-1} f_m(\rho) | \mathbb{1}, m \rangle \\ &= (1-r) \langle K, m | (1-A)^{-1} \rho f_m(\rho) | \mathbb{1}, m \rangle \\ &= (1-r) \sum_{l=-\infty}^K r^{-l} f_m(r^{-l}) , \end{aligned} \quad (49)$$

where the derivative ∂_ρ is defined as in (25)

$$\partial_\rho := \frac{1}{(1-r)\rho} (1-A) .$$

The last expression for the integral in (49) has the form of a usual Jackson integral [8].

To construct the action, note that the two-dimensional space with the coordinates $z, \bar{z}$ and the symmetry transformations (46) can be considered as a result of the conformal map

$$u \rightarrow z = e^u , \quad \bar{u} \rightarrow \bar{z} = e^{\bar{u}} \quad (50)$$

of a cylinder with the coordinates $\bar{u}, u$. This map is well known and is used widely, in particular, in conformal and string theories (in frame of so called radial quantization, see e.g. [18]). A coordinate along the cylinder is associated with a time coordinate and space coordinate takes values on a circle. After the conformal mapping the coordinate ρ plays the role of time and Φ of space coordinate. It is easy to see that in coordinates $u, \bar{u}$ the transformations (46) becomes translations and the time coordinate $\tau := Re u$ takes values on an equidistant lattice with a spacing $\ln r$. In continuous case a field theory free action for a scalar field Ψ in polar coordinates on the z -plane has the form

$$S_0 = \int_0^\infty \frac{d\rho}{\rho} d\phi \Psi(\rho, \phi) (\rho \partial_\rho \rho \partial_\rho + \partial_\Phi^2) \Psi(\rho, \phi) . \quad (51)$$

This expression is explicitly invariant with respect to dilatations of ρ and translations of ϕ (continuous analog of (46)).

A quantum plane analog has the following form

$$S_0^r = \lim_{K \rightarrow \infty} \sum_{m=-\infty}^{\infty} \langle K, m | (1 - A)^{-1} \Psi_{-m}(\rho) G^{-1} \Psi_m(\rho) | \mathbb{1}, m \rangle .$$

Here the operator G^{-1} has the form similar to the Hamiltonian H

$$G^{-1} = \frac{H}{\Omega} ,$$

though its meaning is quite different, of course. The finite difference operator G^{-1} in this action has the eigenvalues $\lambda_{P,m}$ presented in (48). In the case of interacting Ψ^4 theory a one-loop correction to, for example, mass term is proportional to a trace of the operator G

$$\sum_{m=-\infty}^{\infty} \int_0^{-\pi/\ln r} \frac{dP}{\lambda_{P,m}} ,$$

which has no divergencies. Note, however, that summation over orbital number m is the same as in continuous case.

5 Conclusion

The main result of this work are the relations (4) and (9) which show that twists of quantum groups and corresponding q-spaces can be realized with help of the auxiliary non-commuting elements e^i, g_k satisfying the relations (5) and (6). Geometrically this can be interpreted as a transition to another coordinate frame on a q-space and so for many problems twisted quantum groups must be physically equivalent. Let us mention again the analogy between the algebra of e^i, g_k and well known quaternions σ_i : one can use either usual three-dimensional coordinates x^i or non-commutative quaternions $\hat{x}^i := x^i \sigma_i$ (no summation). Physical content of a problem is not changed. Unfortunately, many important quantum groups, for example, orthogonal ones, have not twisted counterparts with commuting coordinates of a corresponding quantum space. Note, however, that linear groups play important role as groups of space-time symmetries. It is enough to mention $SL(2, C)$ as universal covering of Lorentz group and $SU(2, 2)$ as covering of four-dimensional conformal group.

Using q-deformed space with commuting coordinates we constructed multidimensional q-group invariant generalization of the famous Jackson derivatives.

As another application of q-spaces with commuting coordinates we considered quantum mechanics and simple field theory on a two-dimensional quantum space. The structure of the involution leads to the mixed coordinate-angular momentum representation of states of the system. This, in turn, results in discretization of only one (radial) coordinate of the space in spite of the q-deformed differential calculus,

the spectrum of the angular momentum operator being unbounded as in usual continuous case. Such partial discretization of a space-time is enough for two-dimensional models but can be dangerous for higher dimensional cases. Another lesson from the considered models is that q -derivatives or finite differences constructed with help of operators of the form (23) are connected with dilatations and not with translations. This implies that corresponding coordinates are related with usual ones (in which derivatives generate translations) by non-linear exponent-like map of the type (50). It is clear, that symmetry transformations, e.g. of Lorentz group, have quite different form in non-linearly transformed coordinate frames. This remark can be important for numerous attempts to construct quantum deformations of relativistic symmetry.

References

- [1] H.J.Rothe, Lattice Gauge Theories. An Introduction, World Scientific, Singapore, 1992.
- [2] D.Blokhintsev, Space and Time in Microworld, Riedel, Dortrecht, 1973.
- [3] V.G.Drinfel'd, Proc.of the Berkley Inst. Congress of Math. 1 (1987) 798.
- [4] S.L.Woronowicz, Comm.Math.Phys., 111 (1987) 613.
- [5] L.D.Faddeev, N.Yu.Reshetikhin and L.A.Takhtadjan, Algebra i Analis, 1 (1989) 178 (Transl.: Leningrad Math. Jour., 1 (1990) 193).
- [6] J.Wess and B.Zumino, Nucl.Phys.Suppl., 18B (1990) 302.
- [7] F.N. Jackson, Quant. J. Pure and Appl. Math., 41 (1910) 193.
- [8] H.Exton, q -Hypergeometric functions and applications, Ellis Horwood, Chichester, 1983.
- [9] A.Dimakis and F.Müller-Hoissen, Phys.Lett., B295 (1992) 242.
- [10] A. Kempf, Quantum field theory with nonzero minimal uncertainties in positions and momenta, preprint DAMTP-94-33, 1994.
- [11] M.Chaichian and A.Demichev, Jour.Math.Phys., 36 (1995) 398.
- [12] V.G.Drinfel'd, Leningrad Math.Jour., 1 (1989) 1419.
- [13] N.Reshetikhin, Lett.Math,Phys., 20 (1990) 331.
- [14] A.Schirrmacher, Z.Phys., C50 (1991) 321.

- [15] B.Zumino, in Proc. of X-th IAMP Conference, Leipzig, 1991 (preprint LBL-31432, UCB-PTH-62/91).
- [16] A.Schirrmacher, Jour. of Phys., A24 (1991) L1249.
- [17] A.Kempf and S.Majid, Jour.Math.Phys., 35 (1994) 6802.
- [18] M.Green, D.Scharz and E.Witten, Superstring Theory, v.1, Cambridge Univ. Press, Cambridge, 1987.